

2. Серонегативний щодо кашлюку прошарок дорослого населення становить 50%, що свідчить про необхідність проведення імунопрофілактики дорослому населенню.

3. Можливість повторної ревакцинації проти кашлюку дорослих осіб на даний час в Україні відсутня.

4. У Львівській області реєструються поодинокі випадки захворювання на кашлюк серед дорослих осіб при недостатній настороженості медичних працівників щодо захворюваності на дану нозологію дорослого населення, що свідчить про необхідність

здійснювати лабораторні обстеження на кашлюк осіб з кашлем, що триває понад 14 днів.

5. Відсутність в районних центрах Львівщини можливості обстеження методом ПЛР на наявність *Bordetella pertussis* та методом імуноферментного аналізу для виявлення імуноглобулінів зменшує можливість реєстрації випадків захворювання на кашлюк.

6. Своєчасне виявлення хворих на кашлюк та їх лікування, забезпечить зниження захворюваності на кашлюк населення Львівщини.



ЛІТЕРАТУРА

1. Дуда О.К. Кашлюк і паракашлюк: сучасні підходи до діагностики та лікування (клінічна лекція) / О.К.Дуда // «Мистецтво лікування». - №3-4 – 2016. – С.129-130.
2. Наказ МОЗ України від 15.04.05 №169 «Про затвердження методичних вказівок з мікробіологічної діагностики кашлюку та паракашлюку».
3. Дитячі інфекційні хвороби та перспективи сучасної вакцинології (за матеріалами 5-го Світового конгресу Міжнародного товариства з дитячих інфекційних хвороб) / В.І. Задорожна // Профілактична медицина. - 2008. - № 2. - С. 63–68.
4. Лاپій Ф.І. Актуальность эффективной защиты против коклюша / Ф.І. Лاپій // «Здоровье ребенка». - №3. - 2010. - С. 84-86.

REFERENCES:

1. Duda O.K. Kashlyuk i parakashlyuk: suchasni pidhodi do diagnostiki ta likuvannya (klinichna lekcija) / O.K.Duda // «Mistectvo likuvannya». - №3-4 – 2016. – S.129-130.
2. Nakaz MOZ Ukraini vid 15.04.05 №169 «Pro zatverdzhennya metodichnih vkazivok z mikrobiologichnoi diagnostiki kashlyuku ta parakashlyuku».
3. Dityachi infektsijni hvorobi ta perspektivi suchasnoi vakcinologii (za materialami 5-go Svitovogo kongresu Mizhnarodnogo tovaristva z dityachih infektsijnih hvorob) / V.I. Zadorozhna // Profilaktichna medicina. - 2008. - № 2. - S. 63–68.
3. Lapij F.I. Aktual'nost' effektivnoj zashchity protiv koklyusha / F.I. Lapij // «Zdorov'e rebenka». - №3. - 2010. - S. 84-86.

UDC: 616.379-008.64-002.3-092:612.115

THE STUDY OF CHANGES IN THE BLOOD OF COMPLICATIONS OF DIABETES

Karateieva S.Y.

MD, PhD, Assoc. Professor,

Department of Nursing and Higher Nursing Education

Higher educational establishment Ukraine state «Bukovinian State Medical University», Ukraine.

E-mail: Karatsveta@gmail.com

Golovachuk O.K.

MD, PhD, Assoc. Professor,

Women's counseling № 1 Chernivtsi Ukraine

Berezova M.S.

MD, PhD, Assoc. Professor,

Department of Internal Medicine and Infectious

Higher educational establishment Ukraine state «Bukovinian State Medical University», Ukraine.

E-mail: myroslava@bukovynaonline.com

Cherevuk A.A.

Student of Higher educational establishment Ukraine state «Bukovinian State Medical University», Ukraine.

Karateieva A.O.

Student of Yuriy Fedkovych Chernivtsi National University Chernivtsi. Ukraine

E-mail: nastyakarateieva@gmail.com



Summary. Study in rats experiment rats with pyoinflammatory processes against the background of diabetes using ozone therapy revealed improvement of the regulation indices of blood aggregation state, decrease of hypercoagulation manifestations: increase of time of plasma recalcification, activated partial thromboplastin time, prothrombin time, thrombin time, activation of thrombocyte-vascular hemostasis; significant protective properties with decreased azoalbumin lysis, total nonenzymatic fibrinolytic activity and proteinase activity by Kunitz in blood plasma are detected.



Key words: diabetes, pyoinflammatory complications, ozone therapy.



Introduction. Diabetes topicality is caused by the fact that in more than 30-70% of patients pyoinflammatory processes develop and 50% of hospitalized patients need surgical care for purulent-necrotic complications [2, 6, 8]. The rapid progression of purulent-necrotic foci is caused by the development of polymorphic aerobic and anaerobic non-clostridial infection, which is often irresponsive to most antibiotics [3, 5, 7, 9]. Purulent infection and diabetes complicate the course of one another. Mortality caused by DM with purulent surgical infection, as well as a few decades ago, remains high – from 6 to 44.4% [1, 4, 10, 11, 12].

Thus, pyoinflammatory complications of soft tissues in patients with diabetes are a complex surgical problem. That is why the course of pyoinflammatory processes of soft tissues in diabetic patients requires further study to optimize its correction.

Material and methods. The experimental study was conducted on 35 white mature rats. The control group numbers 8. Diabetes was simulated by subcutaneous administration of alloxan (100mg per kg of body weight). Rats of the first experimental group (10) received subcutaneous administration of fecal suspension (pool of 20 animals' feces was diluted with 0.9% sodium chloride solution) at a dose of 0.5ml per 100g of body weight. On the 14th day after alloxan administration the animals of the second experimental group (9) were injected 10% feces suspension subcutaneously. The animals of the third group (8) with simulated DM were administered fecal mixture subcutaneously and during the next three days ozonized 0.9% sodium chloride solution was administered intraperitoneally once a day at the rate of 1.0ml per 100g of body weight.

Ozonation of sterile 0.9% sodium chloride solution was carried out by the apparatus "Boson" until the concentration of ozone in isotonic sodium chloride solution was 20 mg/ml.

Discussion of the results of the study. Assessment of the system of unlimited proteolysis, fibrinolysis activity and average weight molecules of blood plasma under conditions of purulent processes of skin and subcutaneous tissue in mature rats revealed increase of azocasein lysis and average weight molecules concentration in blood plasma (Table 1).

Table 1. State of system of unlimited proteolysis, fibrinolysis activity and average weight molecules of plasma under conditions of pyoinflammatory processes of soft tissues in mature rats ($x \pm Sx$)

Indices	Control (n=8)	Main Fecal suspension administration (n=10)
Azoalbumin lysis, mg/ml hour	2,24±0,351	2,39±0,329
Azocasein lysis, mg/ml hour	1,87±0,227	3,00±0,209 p<0,01
Azocollagen lysis, mg/ml hour	0,77±0,137	1,14±0,176 p<0,01
Total fibrinolytic activity, mg/ml hour	4,23±0,486	4,78±0,489 p<0,05
Nonenzymatic fibrinolytic activity, mg/ml hour	1,88±1,05	1,11±0,175 p<0,01
Proteinase activity by Kunitz – caseinolytic units, calibrated by trypsin	0,19±0,029	0,27±0,031 p<0,01
Average weight molecules concentration, arbitrary units	0,12±0,015	0,18±0,018 p<0,05

p – probability of differences compared with control,
n – number of observations.

Assessment of regulation system of aggregation state of blood under conditions of purulent processes development in mature rats revealed (Table 2) decrease of time of plasma recalcification, thrombin time, increase of adhesive thrombocytes percentage and index of platelets spontaneous aggregation, indicating activation of platelet-vascular hemostasis and coagulation system of plasma.

Decreasing of time of Hageman-factor-dependent fibrinolysis was ineffective in opposition to the specified changes of hypercoagulation activation.

Diabetes significantly complicated the course of purulent processes. That was indicated by a significant increase of proteolysis by azoalbumin

Table 2. State of regulation system of blood aggregation state under conditions of purulent processes of soft tissues in mature rats ($x \pm Sx$)

Indices	Control (n=8)	Main Fecal suspension administration (n=10)
Time of plasma recalcification, sec	101,5±2,38	90,8±2,33 p<0,01
Activated partial thromboplastin time, sec	42,5±2,32	35,8±2,35 p<0,01
Prothrombin time, sec	16,4±1,38	13,2±1,37 p<0,05
Thrombin time, sec	14,5±0,97	10,9±1,07 p<0,05
Concentration of fibrinogen in blood plasma, g/l	4,48±0,179	4,80±0,175 p<0,05
Antithrombin activity III, %	98,2±3,53	104,6±3,37
Activity of blood clotting XIII factor, %	94,9±3,74	96,5±3,35
Percentage of adhesive thrombocytes by Mishcheko method, %	2,69±0,362	4,90±0,358 p<0,001
Index of thrombocytes spontaneous aggregation by Tacolla method, %	28,3±1,91	34,4±1,90 p<0,05
Hageman-factor-dependent fibrinolysis, min	16,9±0,68	13,4±0,66 p<0,01
Plasminogen potential activity, min	15,8±1,26	12,4±1,19 p<0,01
Antiplasmins, %	93,0±4,75	100,6±4,48 p<0,05

Table 3. State of system of unlimited proteolysis, fibrinolysis activity and concentration of average weight molecules in blood plasma in condition of pyoinflammatory processes in soft tissues in mature rats with diabetes against the background of ozone therapy ($x \pm Sx$)

Indices	Fecal suspension administration with DM (n=9)	Fecal suspension administration with DM against the background of ozone therapy (n=8)
Azoalbumin lysis, mg/ml hour	4,64±0,443	3,08±0,282 p<0,02
Azocasein lysis, mg/ml hour	5,95±0,744	4,41±0,735 p<0,01
Azocollagen lysis, mg/ml hour	3,57±0,743	1,84±0,351 p<0,01
Total fibrinolytic activity, mg/ml hour	9,84±0,838	6,34±0,709 p<0,01
Nonenzymatic fibrinolytic activity, mg/ml hour	7,55±1,495	1,42±0,198 p<0,01
Proteinase activity by Kunitz – caseinolytic units, calibrated by trypsin	0,89±0,103	0,31±0,142 p<0,01
Average weight molecules concentration, arbitrary units	0,30±0,034	0,32±0,039

Table 4. State of regulation system of blood aggregation state under conditions of pyoinflammatory processes in soft tissues in diabetic mature rats against the background of ozone therapy ($x \pm Sx$)

Indices	Feces suspension administration (n=9)	Feces suspension administration using ozone therapy (n=8)
Plasma recalcification time, sec	67,7±4,39	84,8±2,01 p<0,01
Activated partial thromboplastin time, sec	21,3±1,73	37,1±2,39 p<0,001
Prothrombin time, sec	9,31±0,869	14,0±1,38 p<0,01
Thrombin time, sec	7,11±0,579	10,9±0,77 p<0,01
Fibrinogen concentration in blood plasma, g/l	2,99±0,161	3,55±0,225 p<0,05
Antithrombin activity III, %	99,1±6,90	116,3±5,35 p<0,05
Concentration of soluble fibrin monomer complexes in blood, mg/l	3,25±0,403	7,57±0,693 p<0,001
Activity of blood clotting XIII factor, %	88,8±7,61	91,9±5,48 p<0,05
Percentage of adhesive thrombocytes by Mishchenko method, %	11,95±0,558	8,60±0,687 p<0,01
Index of thrombocytes spontaneous aggregation by Tacolla method, %	45,7±2,07	33,8±2,46 p<0,01
Hageman-factor-dependent fibrinolysis, min	9,67±0,758	11,8±1,06 p<0,01
Plasminogen potential activity, min	10,3±0,69	8,33±0,810 p<0,01
Antiplasmins, %	112,9±5,36	115,9±5,10

lysis, azocasein, azocollagen, increased total nonenzymatic fibrinolytic activity and proteinase activity by Kunitz (Table 3). Increase of average weight molecules indicated growth of the intoxication degree. Diabetes also caused significant disorders in hemostasis system under the conditions and presence of purulent processes. Thus, a significant increase of blood plasma hypercoagulation was revealed by the decrease of time of plasma recalcification, activated partial thromboplastin time, prothrombin time, thrombin time. Decrease of fibrinogen concentration in blood plasma denoted its use in the hypercoagulation process. Activation of thrombocyte-vascular hemostasis under these conditions was confirmed by the increase of adhesive thrombocytes percentage and index of platelets spontaneous aggregation.

The use of ozone therapy in conditions of pyoinflammatory processes in mature rats with diabetes discovered significant protective properties, indicated by azoalbumin lysis decrease, total nonenzymatic fibrinolytic activity and proteinase activity by Kunitz in blood plasma (Table 3).

Ozone therapy also improved the indices of regulation of blood aggregation state under conditions of purulent processes presence in diabetic mature rats. This was detected by the decrease of hypercoagulation manifestations, indication of which was an increase of plasma

recalcification time, activated partial thromboplastin time, prothrombin time, thrombin time (Table 4). The decrease of pathogenic influence of thrombocyte-vascular hemostasis activation indicated the decrease of adhesive thrombocytes percentage by Mishchenko and index of thrombocyte spontaneous aggregation by Tacolla method. Additional anticoagulating properties under these conditions were revealed by an increase of concentration of soluble fibrin monomer complexes.

Such changes in mature rats can be considered positive in the course of pyoinflammatory processes: decrease of intoxication, increase of protective properties, decrease of hypercoagulation manifestations against the background of chronic forms of diabetes.

Conclusions. 1. The use of ozone therapy under conditions of abscesses of soft tissues in diabetic mature rats reveals significant protective properties with decreased azoalbumin lysis, total nonenzymatic fibrinolytic activity and proteinase activity by Kunitz in blood plasma.

2. Ozone therapy improves the regulation indices of blood aggregation state under conditions of pyoinflammatory processes in diabetic mature rats with decrease of hypercoagulation manifestations: increase of time of plasma recalcification, activated partial thromboplastin time, prothrombin time, thrombin time.



REFERENCES

1. Беляков Н.А. Сахарный диабет как основной компонент патогенеза метаболического синдрома / Н.А.Беляков, С.Ю.Чубриева // Мед. Акад. Ж.- 2009.- Т. 8, № 1.- С. 116-127.
2. Волховская Н.Б. Состояние окислительно-восстановительного равновесия у крыс при разных схемах введения озонированного физиологического раствора /Н.Б.Волховская, О.Е.Колесова, С.Б.Ткаченко //Бюл. экс. биол. и мед.- 2008.- Т. 145, № 2.- С. 141-143.
3. Велигоцкий Н.Н. Динамика морфологических изменений в эпителии раневого канала под воздействием озонотерапии / Н.Н. Велигоцкий, И.Е. Бугаков// Харківська хірургічна школа.-2009.-№ 4.1(36).-С.341-344.
4. Галушко О.А. Гіпоглікемічна та лактацидемічна коми / О.А. Галушко // Ж. практичного лікаря.-2009.-№ 1.-С.24-29.
5. Моравец Тадеуш. Застосування озону в лікуванні утрудненого загоювання після екстраційних ран та деяких патологічних змін слизової оболонки порожнини рота. Початкові спостереження / Тадеуш Моравец, Магдалена Веснер, Ізабелла Коволь [та ін.] // Хірургічна стоматологія.- 2008.- № 1 (54). - С. 86-91.
6. Клімова О.М. Динамика изменения иммунологических показателей у больных после гипнотерапии / О.М.Клімова, П.А.Цай, Т.І.Кордон // Валеологія сучасний стан, напрямки та перспективи розвитку: матеріали VI Міжнародної науково-практичної конференції.- 2008, Т.2. – С. 65-74.
7. Щербатюк Т.Г. Применения озона в медицине: проблема и перспективы / Т.Г.Щербатюк // Фізіол. ж.- 2008.- Т. 54, № 2.- С. 41-48.
8. Ahsen B. Avoiding hypoglycemia: a key to success for glucose – lowering therapy in type 2 diabetes / B. Ahsen // Vase. Health. Risk. Man. – 2013. – Vol. 9. – P. 155-163.
9. American Diabetes Association. Standarts of medical care in diabetes – 2014 //Diabetes Care. – 2014. – Vol. 37. – P. S14-S80.
10. Andrews M.A. Diabetes overtreatment in elderly individuals: risky business in need of better management / Andrews M.A., O'Malley PG. // JAMA. – 2014. – Vol. 311. – P. 2326-2327.
11. Atkinson M.A. Current concepts of the pathogenesis of type 1 diabetes — considerations for attempts to prevent and reverse the disease / Atkinson M.A., Herrath M., Powers A.C., Clare-Salzer M. // Diabetes Care. – 2015. – Vol. 38. – P. 979-988.
12. Bolli G.B. New long-acting insulin analogs: from damp studies to clinical practice / Bolli G.B., De Vries J.H. // Diabetes Care. – 2015. – Vol. 38. – P. 541-543.