



p-ISSN 2224-0721  
e-ISSN 2307-1427

International journal of endocrinology

# МІЖНАРОДНИЙ ЕНДОКРИНОЛОГІЧНИЙ ЖУРНАЛ

Індексується: РІНЦ (Science Index), Index Copernicus International, Google Scholar, Ulrichsweb Global Serials Directory, «Джерело», Academic Resource Index (Research Bible)


ДЕСЯТЬ РОКІВ  
РАЗОМ з читацями ...

[www.mif-ua.com](http://www.mif-ua.com)

5(69) • 2015

СПЕЦІАЛІЗОВАНИЙ РЕЦЕНЗОВАНИЙ  
НАУКОВО-ПРАКТИЧНИЙ ЖУРНАЛ

ЗАСЛАВСКИЙ  
Издательский дом



# Міжнародний ендокринологічний журнал

# Международный эндокринологический журнал

# International journal of endocrinology

Спеціалізований рецензований науково-практичний журнал

Засновано у вересні 2005 року

Періодичність виходу: 8 разів на рік

Журнал включений до наукометричних та спеціалізованих баз даних РІНЦ (Science Index), Index Copernicus International, Google Scholar,

Ulrichsweb Global Serials Directory, «Джерело»,

Academic Resource Index (Research Bible)

Імпакт-фактор РІНЦ 0,064

№ 5(69), 2015





**Міжнародний  
ендокринологічний  
журнал**

*Міжнародний спеціалізований  
рецензований  
науково-практичний журнал*

*Міжнародний специалізованний  
рецензований  
науково-практичний журнал*

*International specialized reviewed  
practical-scientific journal of endocrinology*

*Журнал включений до наукометрических  
та спеціалізованих баз даних РІНЦ (Science Index),  
Index Copernicus International, Google Scholar,  
Ulrichsweb Global Serials Directory, «Джерело»,  
Academic Resource Index (Research Bible)*  
*Імпакт-фактор РІНЦ 0,064*

**№ 5(69) • 2015**

**ISSN 2224-0721**

**Передплатний індекс: 94553**



**Співзасновники:**

**Буковинський державний медичний  
університет,  
Заславський О.Ю.**

**Шеф-редактор Заславський О.Ю.**

**Завідуюча редакцією Брандіс Т.Я.**

**Заступник завідуючої редакцією**

**Курпріченко Н.В.**

**Адреса для звертання:**

**Із питань передплати:** info@mif-ua.com,  
тел. +38 (044) 223-27-42

**З питань розміщення реклами  
та інформації про лікарські засоби:**

reclama@mif-ua.com,  
office@zaslavsky.kiev.ua  
golubinichayan@gmail.com

*Журнал внесено до переліку наукових  
фахових видань України. Наказ МОН України  
від 15.04.2014 р. № 455*

*Рекомендуються до друку та до поширення  
через мережу Інтернет вченого радио  
Вищого державного навчального закладу  
IV рівня акредитації «Буковинський державний  
медичний університет» МОЗ України  
(27 серпня 2015 р., протокол № 1).*

**Російською, українською та англійською мовами**  
Реєстраційне посвідчення  
КВ № 19313-9113ПР.

Видано Державною реєстраційною  
службою України 06.09.2012 р.

Формат: 60×84/8. Ум. друк. арк. 21,86  
Тираж 3 000 прим.

**Адреса редакції:**

а/с 74, м. Київ, 04107, Україна  
Тел./факс: +38(044) 223-27-42

E-mail: medredactor@i.ua

(Тема: До редакції «Міжнародного  
ендокринологічного журналу»)  
[www.mif-ua.com](http://www.mif-ua.com)

**Видавець Заславський О.Ю.**

Адреса для листування: а/с 74, м. Київ, 04107  
Адреса реєстрації: пр. Ленінський, 25/126,  
м. Донецьк, 83102

Свідоцтво суб'єкта видавничої справи  
ДК № 2182 від 13.05.2005 р.

**Друкарня «Астро»**

Адреса реєстрації:

вул. Артема, 63а, м. Донецьк, 83001.  
Свідоцтво суб'єкта видавничої справи  
ДК № 2047 від 24.12.2004 р.

Редакція не завжди поділяє думку автора публікації. Відповідальність за вірогідність фактів, власних імен та іншої інформації, використаної в публікації, несе автор. Передрук та інше відтворення в якій-небудь формі в цілому або частково статей, ілюстрацій або інших матеріалів дозволені тільки при попередній письмовій згоді редакції та з обов'язковим посиланням на джерело. Усі права захищені.

**Головний редактор**  
**Володимир Іванович ПАНЬКІВ**

**Науковий редактор**  
**Тарас Миколайович БОЙЧУК**

**Редакційна колегія**

Авраменко Т.В. (Київ)  
Бобирюва Л.Є. (Полтава)  
Боднар П.М. (Київ)  
Большова О.В. (Київ)  
Бондаренко В.О. (Харків)  
Бошюрко В.І. (Івано-Франківськ)  
Вернигородський В.С. (Вінниця)  
Власенко М.В. (Вінниця)  
Волошин О.І. (Чернівці)  
Генделека Г.Ф. (Одеса)  
Гончарова О.А. (Харків)  
Єфімов А.С. (Київ)  
Зелінська Н.Б. (Київ)  
Зубкова С.Т. (Київ)  
Іванов Д.Д. (Київ)  
Іващук О.І. (Чернівці)  
Караченцев Ю.І. (Харків)  
Кирилюк М.Л. (Київ)  
Козаков О.В. (Харків)  
Комісаренко Ю.І. (Київ)  
Корпачов В.В. (Київ)  
Кравченко В.І. (Київ)  
Кравчун Н.О. (Харків)  
Ларін О.С. (Київ)  
Лучицький Є.В. (Київ)  
Макар Р.Д. (Львів)  
Маньківський Б.М. (Київ)  
Марцинік Є.М. (Дніпропетровськ)  
Медведь В.І. (Київ)  
Мелеховець О.К. (Суми)  
Мислицький В.Ф. (Чернівці)  
Мітченко О.І. (Київ)  
Науменко В.Г. (Київ)  
Нетяженко В.З. (Київ)  
Олійник В.А. (Київ)  
Пасечко Н.В. (Тернопіль)  
Пашковська Н.В. (Чернівці)  
Перцева Т.О. (Дніпропетровськ)  
Пиріг Л.А. (Київ)  
Пічкар Й.І. (Ужгород)  
Пішак В.П. (Чернівці)  
Поворознюк В.В. (Київ)  
Полторак В.В. (Харків)  
Прилуцький О.С. (Донецьк)  
Приступок О.М. (Київ)  
Резніков О.Г. (Київ)  
Селіванова К.Ф. (Сімферополь)  
Сергієнко О.О. (Львів)  
Сидорчук Л.П. (Чернівці)  
Сіренко Ю.М. (Київ)  
Скрипник Н.В. (Івано-Франківськ)  
Соколова Л.К. (Київ)  
Томашевський Я.І. (Львів)  
Тронько М.Д. (Київ)  
Хижняк О.О. (Харків)  
Черенко С.М. (Київ)  
Швед М.І. (Тернопіль)  
Шкала Л.В. (Луганськ)  
Юзленко Т.Ю. (Київ)

**Редакційна рада**

Аметов О.С.  
(Москва, Російська Федерація)  
Аристархов В.Г.  
(Рязань, Російська Федерація)  
Базарбекова Р.Б. (Алмати, Казахстан)  
Валеєва Ф.В.  
(Казань, Російська Федерація)  
Вербовой А.Ф.  
(Самара, Російська Федерація)  
Данилова Л.І. (Мінськ, Білорусь)  
Дедов І.І.  
(Москва, Російська Федерація)  
Зельцер М.Ю. (Алмати, Казахстан)  
Ісмаїлов С.І. (Ташкент, Узбекистан)  
Мамедов М.Н.  
(Москва, Російська Федерація)  
Мельниченко Г.А.  
(Москва, Російська Федерація)  
Метревелі Д. (Тбілісі, Грузія)  
Мірзазаде В. (Баку, Азербайджан)  
Мохорт Т.В. (Мінськ, Білорусь)  
Романчишев А.П.  
(Санкт-Петербург, Російська Федерація)  
Свириденко Н.Ю.  
(Москва, Російська Федерація)  
Трошина К.А.  
(Москва, Російська Федерація)  
Шестакова М.В.  
(Москва, Російська Федерація)  
Prof. Dr. Agaç F. (Тірана, Албанія)  
Prof. Alekna V. (Вільнюс, Литва)  
Prof. Czupryniak L. (Лодзь, Польща)  
Prof. Ferranmini E. (Піза, Італія)  
Prof. Mascarenhas M.R.  
(Лісабон, Португалія)  
Assoc. Prof. Mota M. (Крайова, Румунія)  
Prof. Nauman J.A. (Варшава, Польща)  
Dr. Sc. Radzeviciene L. (Каунас, Литва)  
Prof. Stndl E. (Мюнхен, Німеччина)  
Prof. Szabolcs I. (Будапешт, Угорщина)  
Prof. Taton J. (Варшава, Польща)  
Prof. Yki-Järvinen H. (Гельсінкі, Фінляндія)  
Prof. Zgliczynski S. (Варшава, Польща)

**Відповідальний секретар Іван Іванович ПАВЛУНИК**

УДК 616.379-008.64:577.118

RUSNAK I.T.

Higher State Educational Institution «Bukovinian State Medical University», Chernivtsi, Ukraine

## DIABETES MELLITUS AND SOME MICROELEMENTS

**Summary.** This literature review describes the role of the different trace elements in the development, progression and prevention of diabetes mellitus and its complications. The results of the studies of microelements in blood, urine and hair are presented.

**Key words:** diabetes mellitus, trace elements, insulin, chromium, copper, iodine, selenium, manganese, molybdenum, vanadium, zinc.

Diabetes mellitus (DM) is a chronic disease that occurs either when the pancreas does not produce enough insulin or when the body cannot effectively use the insulin it produces [1]. DM leads to high morbidity and mortality throughout the world [2]. The prevalence of DM in the age groups between 20 to 70 years worldwide was estimated to be 8.3 % in 2013 and 10.1 % in 2035. The total number of adult with DM is projected to rise from 382 million in 2013 to 592 million in 2035. In 2013, an estimated 5.1 million people died from consequences of hyperglycemia. Type 2 diabetes mellitus (T2DM) can be prevented or delayed through healthy diet, regular physical activity, maintaining a normal body weight, and avoiding smoking [3].

T2DM is the most common form of DM and individuals are characterized by disorders of insulin action and secretion, either of which may be the predominant feature [4]. Direct associations of trace elements with diabetes mellitus have been observed in many research studies. Insulin action on reducing blood glucose was reported to be potentiated by some trace elements as chromium, vanadium zinc, manganese, molybdenum, and selenium [4].

Trace elements constitute a minute part of the living tissues and have various metabolic characteristics and functions. Trace elements participate in tissue and cellular and subcellular functions; these include immune regulation by humoral and cellular mechanisms, nerve conduction, muscle contractions, membrane potential regulations, and mitochondrial activity and enzyme reactions [4].

The trace element is a dietary mineral that is needed for the proper growth, development, and physiology of the organism. Alterations in the status of trace elements could stem from chronic uncontrolled hyperglycemia. Trace elements include the transition metals vanadium (V), chromium (Cr), manganese (Mn), iron (Fe), cobalt (Co), copper (Cu), zinc (Zn), and molybdenum (Mo) and the nonmetals selenium (Se), fluorine (F), and iodine (I). All of these belong to the category of micro-nutrients, which are needed by the human body in very small quantities (generally less than 100 mg/day) [5].

Chromium is required for normal carbohydrate metabolism and as a critical cofactor for insulin action and is a component of the glucose tolerance factor, which plays a role in glucose homeostasis [4]. Chromium concentrations were significantly reduced in blood of T2DM patients as compared to control subjects of both genders but urinary levels of these elements were found to be higher in the diabetic patients than in the age-matched healthy controls [6].

Copper is considered as both a powerful enzyme catalyst and a dangerous reactant that generates hydroxyl radical. A deficiency of copper results in glucose intolerance, decreased insulin response, and increased glucose response. It is associated with hypercholesterolemia and atherosclerosis [4].

Iodine is absolutely vital for proper thyroid function. The iodine deficiency is the most common cause of hypothyroidism worldwide. The iodine deficiency will cause mental retardation and cretinism and it is the most devastating in all trace elements [4]. Thyroid hormone controls insulin secretion. In hypothyroidism, there is a reduction in glucose-induced insulin secretion by  $\beta$  cells, and the response of  $\beta$  cells to glucose or catecholamine is increased in hyperthyroidism due to increased  $\beta$ -cell mass. Moreover, insulin clearance is increased in thyrotoxicosis [7]. Insulin resistance and  $\beta$ -cell function are inversely correlated with thyroid stimulating hormone (TSH) which may be explained by insulin-antagonistic effects of thyroid hormones along with an increase in TSH. The higher serum TSH usually corresponds to lower thyroid hormones via negative feedback mechanism. As TSH increased, thyroid hormones decreased and insulin antagonistic effects are weakened. These

Адреса для листування з автором:  
Руснак І.Т.  
E-mail: ilona\_svet@mail.ru

© Руснак І.Т., 2015  
© «Міжнародний ендокринологічний журнал», 2015  
© Заславський О.Ю., 2015

observations demonstrate that insulin imbalance is closely associated with thyroid dysfunction and the phenomenon is mediated via  $\beta$ -cell dysfunction [8]. The significantly lower level of iodine was reported in the urine of T2DM than in that of healthy control subjects [9].

Selenium, a trace element, is involved in the complex system of defense against oxidative stress through selenium-dependent glutathione peroxidases and other selenoproteins [4]. Due to its antioxidant properties, selenium might be preventing the development of diabetes. In addition, selenate, an inorganic form of selenium, mimics insulin activity in experimental models [4]. Selenium is incorporated into selenoproteins [10, 11] that have a wide range of pleiotropic effects, ranging from antioxidant and anti-inflammatory effects to the production of active thyroid hormone. In the past 10 years, the discovery of disease-associated polymorphisms in selenoprotein genes has drawn attention to the relevance of selenoproteins to health. Low selenium status has been associated with increased risk of mortality, poor immune function, and cognitive decline. Higher selenium status or selenium supplementation has antiviral effects, is essential for successful male and female reproduction, and reduces the risk of autoimmune thyroid disease [10]. This antioxidant property of selenium prevents the development of complications in diabetic patients [4]. Conflicting evidence linking selenium to glucose metabolism has been reported. For instance, high selenium status was associated with reduced diabetes prevalence in several prospective cohort and case-control studies [12].

Manganese plays an important role in a number of physiologic processes as a constituent of some enzymes such as pyruvate carboxylase and arginase and an activator of different enzymes such as phosphoenolpyruvate carboxykinase and glutamine synthetase. These manganese activated enzymes play important roles in the metabolism of carbohydrates, aminoacids, and cholesterol. Manganese helps in glucose metabolism and it is required for normal synthesis and secretion of insulin. The level of manganese is lower in T2DM subjects as compared to control subjects [4]. In another study, the mean manganese was significantly low in blood and scalp-hair samples of diabetic patients as compared to control and both genders [13].

There is accumulating evidence that molybdenum plays an important role in insulin action as it has been suggested to be able to bind the insulin receptor and to be involved in the activation of glucose metabolism enzymes [11]. Therefore, molybdenum complexes have been proposed as possible adjunct in the treatment of DM [11, 14].

In addition to reproductive problems and skeletal abnormalities observed in case of deficiency, vanadium is likely to have a significant role in thyroid, iron, glucose and lipid metabolism [15]. The potential use of vanadium in the treatment of diabetic complications including cardiomyopathy has been assessed and indeed its hypoglycemic effect along with reversal of functional abnormalities has been clearly demonstrated by several studies [2, 16, 17].

Zinc plays an important role in glucose metabolism. It helps in the utilization of glucose by muscle and fat cells. It is required as a cofactor for the function of intracellular

enzymes that may be involved in protein, lipid, and glucose metabolism. Zinc may be involved in the regulation of insulin receptor-initiated signal transduction mechanism and insulin receptor synthesis [18]. Zinc is a structural part of key antioxidant enzymes such as superoxide dismutase, and zinc deficiency impairs their synthesis, leading to increased oxidative stress. Low zinc has also been seen to lead to poor or slowed wound healing, which is common in diabetic patients. Oxidative stress plays an important role in the pathogenesis of diabetes and its complications. Clinical studies reported that serum levels of zinc are usually found low in T2DM patients compared to nondiabetic due to the impaired intestinal reabsorption of endogenous zinc and the increase in excretion of zinc into the intestine during the digestive process may lead to this low serum zinc level [4].

Zinc supplementation can be effective for preventing or ameliorating of DM. Zinc transporter (ZnT-8) is a crucial protein for the regulation of insulin secretion in pancreatic  $\beta$ -cells [19]. Several studies have reported that diabetics had lower serum/plasma zinc levels [20, 21]. Increased urinary excretion of zinc suggests a deficiency in blood zinc and further dysregulation of insulin secretion [11].

Nutrition management aims to improve health quality maintaining blood glucose levels in normal range so as to reduce the risk for diabetes complications. A well-balanced diet will maintain the impairment of essential macro- and micronutrients in patient with DM.

## Список літератури / References

1. World Health Organization. *Diabetes. Fact Sheet*. — 2013. — (312); <http://www.who.int/mediacentre/factsheets/fs312/en/>
2. Ozturk N. *Trace elements in diabetic cardiomyopathy: An electrophysiological overview* / N. Ozturk, Y. Olgar, S. Ozdemir // *World J. Diabetes*. — 2013. — Vol. 4(4). — P. 92-100. doi: 10.4239/wjd.v4.i4.92
3. International Diabetes Federation. *The IDF Diabetes Atlas*. — 6<sup>th</sup> ed. — Brussels, Belgium: International Diabetes Federation, 2013.
4. Siddiqui K. *Variation in Macro and Trace Elements in Progression of Type 2 Diabetes* / K. Siddiqui, N. Bawazeer, S.S. Joy // *Scientific World Journal*. — 2014. — Vol. 2014. — 461591. doi: 10.1155/2014/461591
5. Fraga C.G. *Relevance, essentiality and toxicity of trace elements in human health* // *Molecular Aspects of Medicine*. — 2005. — Vol. 26(4–5). — P. 235-244.
6. Kazi T.G., Afridi H.I., Kazi N. et al. *Copper, chromium, manganese, iron, nickel, and zinc levels in biological samples of diabetes mellitus patients* // *Biological Trace Element Research*. — 2008. — Vol. 122(1). — P. 1-18.
7. Mitrou P., Raptis S.A., Dimitriadis G. *Insulin action in hyperthyroidism: a focus on muscle and adipose tissue* // *Endocrine Reviews*. — 2010. — Vol. 31(5). — P. 663-679. doi: 10.1210/er.2009-0046
8. Chen G., Wu J., Lin Y. et al. *Associations between cardiovascular risk, insulin resistance,  $\beta$ -cell function and thyroid dysfunction: a cross-sectional study in She ethnic minority group of Fujian Province in China* // *European Journal of Endocrinology*. — 2010. — Vol. 163(5). — P. 775-782. doi: 10.1530/EJE-10-0710

9. Al-Attas O.S., Al-Daghri N.M., Alkharfy K.M. et al. Urinary iodine is associated with insulin resistance in subjects with diabetes mellitus type 2 // Experimental and Clinical Endocrinology and Diabetes. — 2012. — Vol. 120(10). — P. 618-622. doi: 10.1055/s-0032-1323816
10. Rayman M.P. Selenium and human health / M. P. Rayman // The Lancet. — 2012. — Vol. 379, № 9822. — P. 1256-1268. Doi: [http://dx.doi.org/10.1016/S0140-6736\(11\)61452-9](http://dx.doi.org/10.1016/S0140-6736(11)61452-9)
11. Association of urinary metal profiles with altered glucose levels and diabetes risk: a population-based study in China / Wei Feng, Xiuqing Cui, Bing Liu, Chuanyao Liu [et al.] // PLoS One. — 2015. — Vol. 10(4). — e0123742. doi: 10.1371/journal.pone.0123742
12. Plasma selenium and risk of dysglycemia in an elderly French population: results from the prospective Epidemiology of Vascular Ageing Study / Akbaraly T.N., Arnaud J., Rayman M.P., Hininger-Favier I. [et al.] // Nutr. Metab. (Lond). — 2010. — Vol. 7. — 21. doi: 10.1186/1743-7075-7-21
13. Copper, chromium, manganese, iron, nickel, and zinc levels in biological samples of diabetes mellitus patients / Kazi T.G., Afiridi H.I., Kazi N. [et al.] // Biological Trace Element Research. — 2008. — Vol. 122(1). — P. 1-18. doi: 10.1007/s12011-007-8062-y
14. Comparison of anti-hyperglycemic effect amongst vanadium, molybdenum and other metal maltol complexes / Thompson K.H., Chiles J., Yuen V.G., Tse J. [et al.] // J. Inorg. Biochem. — 2004. — Vol. 98. — P. 683-690.
15. Alternative therapies for diabetes and its cardiac complications: role of vanadium / Clark T.A., Deniset J.F., Heyliger C.E., Pierce G.N. // Heart Fail. Rev. — 2014. — Vol. 19(1). — P. 123-32. doi: 10.1007/s10741-013-9380-0
16. Restoration of cardiomyocyte function in streptozotocin-induced diabetic rats after treatment with vanadate in a tea decoction / Clark T.A., Maddaford T.G., Tappia P.S., Heyliger C.E. [et al.] // Curr. Pharm. Biotechnol. — 2010. — Vol. 11. — P. 906-910.
17. A tea/vanadate decoction delivered orally over 14 months to diabetic rats induces long-term glycemic stability without organ toxicity / Clark T.A., Heyliger C.E., Kopilas M., Edel A.L. [et al.] // Metabolism. — 2012. — Vol. 61. — P. 742-753. doi: 10.1016/j.metabol.2011.10.007
18. Tang X., Shay N.F. Zinc has an insulin-like effect on glucose transport mediated by phosphoinositol-3-kinase and Akt in 3T3-L1 fibroblasts and adipocytes // Journal of Nutrition. — 2001. — Vol. 131(5). — P. 1414-1420.
19. Identification and cloning of a beta-cell-specific zinc transporter, ZnT-8, localized into insulin secretory granules / Chimienti F., Devergnas S., Favier A., Seve M. // Diabetes. — 2004. — Vol. 53. — P. 2330-2337.
20. Plasma mineral content in type-2 diabetic patients and their association with the metabolic syndrome / Aguilar M.V., Saavedra P., Arrieta F.J., Mateos C.J. [et al.] // Ann. Nutr. Metab. — 2007. — Vol. 51. — P. 402-406.
21. Ferdousi S., Mia A.R. Serum levels of copper and zinc in newly diagnosed type-2 diabetic subjects // Mymensingh Med. J. — 2012. — Vol. 21. — P. 475-478.

**Отримано 01.10.15** ■

Руснак І.Т.

Вищий державний навчальний заклад «Буковинський державний медичний університет»,  
м. Чернівці

### ЦУКРОВИЙ ДІАБЕТ ТА ДЕЯКІ МІКРОЕЛЕМЕНТИ

**Резюме.** В огляді літератури описана роль різних мікроелементів у розвитку, прогресуванні та профілактиці цукрового діабету та його ускладнень. Наведено результати досліджень мікроелементів у крові, сечі, волоссі.

**Ключові слова:** цукровий діабет, мікроелементи, інсулін, хром, мідь, йод, селен, марганець, молібден, ванадій, цинк.

Руснак И.Т.

Высшее государственное учебное заведение  
«Буковинский государственный медицинский университет», г. Черновцы

### САХАРНЫЙ ДИАБЕТ И НЕКОТОРЫЕ МИКРОЭЛЕМЕНТЫ

**Резюме.** В обзоре литературы описана роль разных микроэлементов в развитии, прогрессировании и профилактике сахарного диабета и его осложнений. Приведены результаты исследований микроэлементов в крови, моче, волосах.

**Ключевые слова:** сахарный диабет, микроэлементы, инсулин, хром, медь, йод, селен, марганец, молибден, ванадий, цинк.